arXiv:0906.0428v1 [math.PR] 2 Jun 2009

LARGE DEVIATIONS OF U-EMPIRICAL KOLMOGOROV-SMIRNOV TESTS, AND THEIR EFFICIENCY

Yakov Nikitin [1]

St.Petersburg State University

Non-degenerate U-empirical Kolmogorov-Smirnov tests are studied and their large deviation asymptotics under the null-hypothesis is described. Several examples of such statistics used for testing goodness-of-fit and symmetry are considered. It is shown how to calculate their local Bahadur efficiency.

1 Introduction.

Let $X_1, X_2, ...$ be i.i.d. observations with continuous distribution function (df) F. Denote by F_n the usual empirical df (edf) based on the first n observations. The Kolmogorov statistic is the distribution-free statistic defined by

$$D_n = \sup_t |F_n(t) - F(t)|. \tag{1}$$

In order to describe the large deviation asymptotics of statistic (1) consider for $0 < a < 1$ the function

$$f(a,t) = \begin{cases} (a+t)\,\ln\dfrac{a+t}{t} + (1-a-t)\,\ln\dfrac{1-a-t}{1-t}\,, & 0 \le t \le 1-a\,, \\ \\ +\infty\,, & 1-a < t \le 1\,, \end{cases}$$

and put

$$f_0(a) = \inf_{0<t<1} f(a,t)\,.$$

The following theorem was proved in [1], see also [2] and [3].

Theorem 1.1. *For any $a \in (0,1)$ we have*

$$\lim_{n\to\infty} n^{-1} \ln \mathbb{P}(D_n > a) = -f_0(a),$$

---

[1] Partially supported by RFBR grant No. 07-01-00159, and by grant NSh. 638.2008.1.

where the function $f_0(a)$ *is continuous on* $(0,1)$, *and as* $a \to 0$

$$f_0(a) = 2a^2(1+o(1))\,.$$

In particular, this result can be used to calculate Bahadur efficiency of various modifications of Kolmogorov-Smirnov tests, see [2] and [3].

We are interested in U-empirical generalizations of Theorem 1.1 and their applications to the problem of testing goodness-of-fit and symmetry. Let $h(x_1, ..., x_m)$ be a real-valued symmetric kernel of degree $m \geq 1$. Consider the U-empirical df (udf)

$$G_n(t) = \binom{n}{m}^{-1} \sum_{1 \leq i_1 < ... < i_m \leq n} I\{h(X_{i_1}, ..., X_{i_m}) < t\}, \quad t \in R^1.$$

The properties of such udf's and their use in Statistics were studied in [4, 5, 6, 7]. Also denote

$$G(t) = \mathbb{P}(h(X_1, ..., X_m) < t) \tag{2}$$

and assume that this df is continuous and (for simplicity) strictly monotonic. Then the U-empirical analog of the Kolmogorov statistic (1) has the form

$$DU_n = \sup_t |G_n(t) - G(t)|$$

and coincides with the Kolmogorov statistic when $m = 1$ and $h(t) = t$.

Various tests of goodness-of-fit and symmetry can be constructed utilizing the Kolmogorov-Smirnov-type statistic

$$SU_n = \sup_t |G_n(t) - F_n(t)|, \tag{3}$$

with various choices of the kernel h. Statistics of this type frequently appear in constructing statistical tests based on characterizations of distributions; then it is assumed that $G(t) \equiv F(t)$, see [8, 9, 10, 11].

The next example illustrates the idea of building such tests. Consider scale-free testing of exponentiality based on Desu's characterization [12]: *Let* X_1 *and* X_2 *be independent non-degenerate and non-negative rv's with common df* F. *Then the rv's* $2\min(X_1, X_2)$ *and* X_1 *have the same distribution if and only if* $F(x) = 1 - \exp(-\lambda x), x \geq 0$ *for some* $\lambda > 0$.

Let us compare the U-empirical df

$$\tilde{G}_n(t) = \binom{n}{2}^{-1} \sum_{1 \leq j < k \leq n} I\{2\min(X_j, X_k) < t\}$$

with the usual edf $F_n(t)$ by considering the statistic

$$DE_n = \sup_t |\tilde{G}_n(t) - F_n(t)|$$

and assuming its large values to be critical. The limiting distribution of this statistic can be found using the results of [13] where weak convergence of U- empirical processes is studied, see also [14] and [4]. Critical values of DE_n can be calculated via simulation.

The logarithmic large deviation asymptotics of DE_n enables to calculate its Bahadur efficiency and compare it with other statistics used for testing exponentiality, see [2, 3, 15]. Various examples of similar statistics will be given at the end of this paper.

The problem of large deviation asymptotics for U-statistics is studied insufficiently. The abstract large deviation principle stated in [16] and [7] is non-effective for statistical applications. The point is that the rate of decrease for large deviation probabilities is given implicitly, as a value of complicated extremal problem in which the Kullback-Leibler information is minimized on an involved set of probability measures. The result that could actually be applied to the efficiency calculations was first obtained for bounded kernels in [18]. This result is stated as follows.

Theorem 1.2. *Consider the U-statistic of degree $m \geq 1$*

$$U_n = \binom{n}{m}^{-1} \sum_{1 \leq i_1 < \ldots < i_m \leq n} \Phi(X_{i_1}, ..., X_{i_m})$$

with centred, bounded, and non-degenerate real-valued kernel Φ, so that

$$E\Phi(x_1, \ldots, X_m) = 0, \quad |\Phi(x_1, ..., x_m)| \leq M,$$

and $\sigma^2 = E\varphi^2(X_1) > 0$ with $\varphi(y) = E(\Phi(X_1, ..., X_m)|X_1 = y)$. Then we have

$$\lim_{n\to\infty} n^{-1} \ln \mathbb{P}(U_n \geq a) := -g(a|\Phi) = -\sum_{j=2}^{\infty} b_j a^j, \tag{4}$$

where the series converges for sufficiently small $a > 0$ and $b_2 = (2m^2\sigma^2)^{-1}$.

The aim of the present paper is to extend Theorem 1.1 (using Theorem 1.2) to the case of statistics DU_n, SU_n, their one-sided variants, and some other similar statistics. In the last section we show how to use the large deviation results for calculating local Bahadur efficiencies.

2 Auxiliary arguments and formulation of results

To study large deviations of Kolmogorov-type statistics, it is natural to begin with a simple one-sided statistics, see [1], [2],

$$DU_n^+ = \sup_t [G_n(t) - G(t)], \quad DU_n^- = \sup_t [G(t) - G_n(t)].$$

For any $\varepsilon > 0$ denote $P_n(\varepsilon) = \mathbb{P}(DU_n \geq \varepsilon)$, and let $P_n^+(\varepsilon)$ and $P_n^-(\varepsilon)$ be the corresponding probabilities for DU_n^+ and DU_n^-. It is evident that

$$\max(P_n^+(\varepsilon),\ P_n^-(\varepsilon)) \leq\ P_n(\varepsilon) \leq\ 2\max(P_n^+(\varepsilon), P_n^-(\varepsilon)). \tag{5}$$

Therefore, if we prove that for some functions g^+ and g^-

$$\lim_{n\to\infty} n^{-1} \ln P_n^+(\varepsilon) = -g^+(\varepsilon), \quad \lim_{n\to\infty} n^{-1} \ln P_n^-(\varepsilon) = -g^-(\varepsilon),$$

then, by (5), we obtain

$$\lim_{n\to\infty} n^{-1} \ln P_n(\varepsilon) = -\min(g^+(\varepsilon), g^-(\varepsilon)). \tag{6}$$

The above argument is also valid for the statistic SU_n when we use one-sided statistics SU_n^+ and SU_n^-. For this reason, below we consider large deviations of one-sided test statistics only.

Let us impose some restrictions on the kernel. We see that the statistic DU_n^+ is, in fact, the supremum of the family of U-statistics with the kernels

$$\Theta(x_1, \dots, x_m; t) = I\{h(x_1, ..., x_m) < t\} - G(t), \quad t \in R^1, \tag{7}$$

depending on t. The same is true for the statistic SU_n^+, where the kernels also depend on t and have the form

$$\Psi(x_1, \dots, x_m; t) = I\{h(x_1, ..., x_m) < t\} - m^{-1} \sum_{i=1}^{m} I\{x_i < t\}, \quad t \in R^1. \tag{8}$$

For any t the kernels $\Theta(\cdot\ ; t)$ and $\Psi(\cdot\ ; t)$ are centred and bounded. Consider their projections

$$\theta(s; t) = E(\Theta(X_1, ..., X_m; t)|X_1 = s), \quad \psi(s; t) = E(\Psi(X_1, ..., X_m; t)|X_1 = s),$$

and the variance functions

$$\sigma_\theta^2(t) = E\theta^2(X_1; t), \quad \sigma_\psi^2(t) = E\psi^2(X_1; t).$$

In the sequel, we will work with general families of kernels $\Phi(\cdot\ ; t \in \mathcal{T})$, that comprise, as particular cases, the families $\Theta(\cdot\ ; t)$ and $\Psi(\cdot\ ; t)$ defined by (7) and (8). For definiteness, we assume that the parameter set $\mathcal{T}$ is some finite or infinite interval $[a, b]$ of the real line.

Most results in the literature related to large deviations of U-statistics deal with U-statistics with specific kernels, see, for example, [17, 16, 18, 19]. Unlike these results, we study whole *families* of kernels and the corresponding *families* of U-statistics indexed by real parameter t. This is a distinctive feature of the paper.

We say that the family of U-statistics $\{U_n(t), t \in [a, b]\}$ with the kernels $\Phi(\cdot\ ; t)$ and projections $\varphi(\cdot\ ; t)$ is *non-degenerate, if its variance function* $\sigma^2_\varphi(t) = E\varphi^2(X_1; t)$ *can vanish only at the ends of the interval* $[a, b]$ *and at a finite number of points in the interior of this interval.*

For example, the one-sided variant DE^+_n of Desu's statistic (1) has the variance function equal to $\frac{1}{4}\exp(-t)(1 - \exp(-t)), t \geq 0$. Another typical variance function that appears below in Example 5 is equal to $\frac{1}{4}t^2(1 - |t|), -1 \leq t \leq 1$. Both families of U-statistics are non-degenerate according to our definition.

Denote by φ^2_0 the maximum of the variance function, i.e., $\varphi^2_0 = \sup_t \sigma^2_\varphi(t) > 0$ and introduce the following subset of the parameter set :

$$T = \{t|\ t \in [a, b],\ \sigma^2_\varphi(t) > \frac{1}{2}\varphi^2_0\}.$$

On this set the variance function is separated from zero. Denote by T^c the complement of T in $[a, b]$.

Consider the family of U-statistics $\{U_n(t), t \in [a, b]\ \}$ corresponding to the family of kernels $\{\Phi(\cdot\ ; t)\}$. For $t \in T$ the kernels are non-degenerate, centred, and bounded. Then the application of Theorem 1.2 yields relation (4) with the function of two arguments

$$g(a, t|\Phi) = \sum_{j=2}^{\infty} b_j(t)a^j \tag{9}$$

where the coefficients $b_j(t)$ depend on t unlike the function $g(a|\Phi)$. Put

$$g_T(a|\Phi) := \inf_{t\in T} g(a, t|\Phi). \tag{10}$$

It turns out that this function determines the large deviation asymptotics of the statistics under consideration, and that the behavior of the function $t \to g(a, t|\Phi)$ on the set T^c is of no importance. The set T can be somewhat enlarged or narrowed without essential changes in the results.

As usually in large deviation theory, deriving the large deviation asymptotics will consist in obtaining upper bound and lower bound. Asymptotic coincidence of the bounds would give the desired result. We begin with the lower bound.

Theorem 2.1. *Suppose that the non-degenerate family of kernels* $\{\Phi(x_1, ..., x_m; t),$ $t \in [a, b]\}$ *is bounded and centred for all* t. *Then for* $a > 0$ *we have*

$$\liminf_{n\to\infty} n^{-1} \ln \mathbb{P}(\sup_t U_n(t) > a) \geq -g_T(a|\Phi). \tag{11}$$

The function $a \to g_T(a|\Phi)$ *is continuous for sufficiently small* a *and admits the representation*

$$g_T(a|\Phi) = \frac{a^2}{2m^2\varphi_0^2} + O(a^3), \;\; a \to 0. \tag{12}$$

From this theorem the lower bound on the large deviation probabilities for U-empirical Kolmogorov-Smirnov tests follows. Both one-sided statistics DU_n^+ and DU_n^- satisfy the conditions of Theorem 2.1. They correspond to different families of kernels $\Theta(\cdot; t)$ and $-\Theta(\cdot; t)$ with common variance function $\sigma_\theta^2(t)$. The corresponding functions $g_T(a|\Theta)$ and $g_T(a| - \Theta)$ are also different, but the first terms of their asymptotic expansions as $a \to 0$ are the same. Now we obtain from (11)

$$\liminf_{n\to\infty} n^{-1} \ln \mathbb{P}(DU_n^\pm > a) \geq -g_T(a| \pm \Theta), \tag{13}$$

and similarly

$$\liminf_{n\to\infty} n^{-1} \ln \mathbb{P}(SU_n^\pm > a) \geq -g_T(a| \pm \Psi). \tag{14}$$

For the purpose of obtaining the upper bound, we assume that the following condition of *monotonicity in parameter* on the family $\{U_n(t), t \in [a, b]\}$ is satisfied. Suppose there exists a sequence of partitions of the interval $[a, b]$ into N parts: $a = t_0 < t_1 < \cdots < t_N = b$, such that the nodes of the partition do not coincide with the zeros of the variance function and that for any $k = 0, \ldots, N - 1$

$$\sup_{t_k \leq t < t_{k+1}} U_n(t) \leq U_n(t_{k+1}) + \Delta_n(N), \tag{15}$$

where the sequence of rv's $\Delta_n(N)$ *decreases fast*, when n and N grow. More precisely, it means that there exists a sequence $\{\tau_N\}, \tau_N \to 0$ as $N \to \infty$, such that

$$\lim_{N\to\infty} \lim_{n\to\infty} n^{-1} \ln \mathbb{P}(\Delta_n(N) > \tau_N) = -\infty. \tag{16}$$

Under this condition the upper bound result takes the following form.

Theorem 2.2. *Suppose that the conditions of Theorem 2.1 are valid and that the condition of monotonicity in parameter of the family* $\{U_n(t)\}$ *holds. Then we have*

$$\limsup_{n\to\infty} \; n^{-1} \ln \mathbb{P}(\sup_t U_n(t) > a) \leq -g_T(a|\Phi). \tag{17}$$

Combining Theorems 2.1 and 2.2, we arrive at the following result.

Theorem 2.3. *Under the conditions of Theorem 2.2 we have*

$$\lim_{n\to\infty} n^{-1} \ln \mathbb{P}(\sup_t U_n(t) > a) = -g_T(a|\Phi).$$

It is shown below that both statistics $DU_n^{\pm}$ and $SU_n^{\pm}$ satisfy the monotonicity condition. If the corresponding families of kernels (7) and (8) are non-degenerate and centred, then Theorem 2.1 is applicable. Hence (5) and (6) imply the following result for the two-sided statistics.

Theorem 2.4. *If the family of kernels* (7) *is non-degenerate, then*

$$\lim_{n\to\infty} n^{-1} \ln \mathbb{P}(DU_n > a) = v_D(a) = -\min(g_T(a|\Theta), g_T(a|-\Theta)).$$

If the family of kernels (8) *is non-degenerate and centred, then*

$$\lim_{n\to\infty} n^{-1} \ln \mathbb{P}(SU_n > a) = v_S(a) = -\min(g_T(a|\Psi), g_T(a|-\Psi)),$$

The functions v_D *and* v_S *are continuous and satisfy the asymptotic relations*

$$v_D(a) \sim -a^2/2m^2\theta_0^2, \quad v_S(a) \sim -a^2/2m^2\psi_0^2, \quad a \to 0.$$

3 Lower bound.

In order to prove the lower bound we need the theorem on implicit analytic operators. Consider three Banach spaces E_1, E_2, and E_3. Denote by $\mathcal{D}_r(x_0, E)$ the ball in the space E of radius r with center at x_0. We are interested in finding solutions $x = x(y)$ of the operator equation

$$F(x, y) = 0 \tag{18}$$

(here y plays a part of a parameter) under the condition

$$x(y_0) = x_0. \tag{19}$$

We assume that the operator $F(x,y)$ is analytic and the condition $F(x_0,y_0)=0$ holds. For main definitions and facts of the theory of analytic operators in Banach spaces, we refer to [20, §22] and [21, §32]. In particular, the operator $F(x,y)$ is called analytic in some domain if in the neighbourhood of any point it can be represented as an uniformly convergent Taylor operator series [20].

Theorem 3.1. *see [20, Theorem 22.2]. Suppose that the operator $F(x,y)$ is analytic in $\mathcal{D}_r(x_0,E_1)\times\mathcal{D}_\rho(y_0,E_2)$ with values in E_3. Let the operator $B=F'_x(x_0,y_0)$ (the derivative is understood in the Fréchet sense) has a bounded inverse operator. Then there exist positive numbers r_1 and ρ_1 such that the equation* (18) *has an unique solution $x=f(y)$ in the ball $\mathcal{D}_{r_1}(x_0,E_1)$. This solution is defined and analytic in the ball $\mathcal{D}_{\rho_1}(y_0,E_2)$, and satisfies the condition* (19)*.*

We will prove Theorem 3.1 using the arguments of [18] in conjunction with some auxiliary results. The main idea is to construct a majorant series for solutions of nonlinear equations that would guarantee the uniform convergence with respect to the parameter t.

Proof of Theorem 3.1. It is clear that for any statistic of the form $\sup_t U_n(t)$, with kernels $\Phi(\cdot\,;t)$ and projections $\varphi(\cdot\,;t)$, the following holds true:

$$\mathbb{P}(\sup_t U_n(t)>a)\geq\sup_{t\in T}\mathbb{P}(U_n(t)>a).$$

Consequently,

$$\liminf_{n\to\infty} n^{-1}\ln\mathbb{P}(\sup_t U_n(t)>a)\geq-\inf_{t\in T} g(a,t|\Phi)=-g_T(a|\Phi).$$

Hence in order to obtain the lower bound we must analyze the function $g(a,t|\Phi)$ in (9) over the set T. Recall that for $t\in T$ we have $\sigma^2_\varphi(t)>\frac{1}{2}\varphi_0^2>0$.

For any Borel set $S\subset R^1$ consider the Banach space of measurable bounded functions $B(S)$ with the norm

$$\parallel x\parallel=\sup\nolimits_{s\in S}|x(s)|\ .$$

We may assume that the initial observations $X_i, i=1,\ldots,n$, are uniformly distributed on $I=[0,1]$. Otherwise we can consider the sample $U_i=F(X_i), i=1,\ldots,n$, and replace the initial kernel $h(X_1,...,X_m)$ by $h(F^{-1}(U_1),\ldots,F^{-1}(U_m))$. The families of kernels $\Phi(\cdot\ ;t),\Psi(\cdot\ ;t)$, $\Theta(\cdot\ ;t)$ and the corresponding families of U-statistics depending on U_i remain centred, bounded, and non-degenerate.

For simplicity we consider only the kernels of degree 2. With slight changes in the proof, the results remain valid for kernels of any finite degree.

For any $t \in T$ we use the variant of Sanov's theorem for large deviations of U-empirical measures from [16], see also [7]. So, we reduce the problem of large deviations to the problem of the minimization of entropy under suitable normalization conditions. It follows from [18] that the function of interest $g(a, t|\Phi)$ is the solution of the extremal problem

$$g(a, t|\Phi) = \inf\{\int_0^1 (1 + ax(s;t)) \ln(1 + ax(s;t))ds : x \in B(I \times T), \int_0^1 x(s;t)ds = 0\} \tag{20}$$

under the normalization condition

$$2\int_0^1 \varphi(s_1;t)x(s_1;t)ds_1 + \int_0^1 \int_0^1 \Phi(s_1, s_2;t)x(s_1;t)x(s_2;t)ds_1 ds_2 = 1. \tag{21}$$

The Euler-Lagrange equation for the extremal $x(s;t)$ from $B(I \times T)$ takes the form, see [18] :

$$\begin{aligned}(1 + ax(s_1;t)) \textstyle\int_0^1 \exp\left\{\lambda(t)a \int_0^1 \Phi(s_1, s_2;t)(1 + ax(s_2;t))ds_2\right\} ds_1 - \\ - \exp\left\{\lambda(t)a \textstyle\int_0^1 \Phi(s_1, s_2;t)(1 + ax(s_2;t))ds_2\right\} = 0,\end{aligned} \tag{22}$$

under the same normalization condition (21). To simplify the notations denote

$$\rho(x|s_1;t) = \int_0^1 \Phi(s_1, s_2;t)x(s_2;t)ds_2.$$

Then (22) can be written as

$$\begin{aligned}(1 + ax(s_1;t)) \textstyle\int_0^1 \exp\left(\lambda(t)a\varphi(s_1;t) + \lambda(t)a^2\rho(x|s_1;t)\right) ds_1 - \\ - \exp\left(\lambda(t)a\varphi(s_1;t) + \lambda(t)a^2\rho(x|s_1;t)\right) = 0,\end{aligned} \tag{23}$$

and the condition (21) becomes

$$\int_0^1 (2\varphi(s_1;t) + a\rho(x|s_1;t))x(s_1;t)ds_1 = 1. \tag{24}$$

Expanding the exponents on the left-hand side of (23) into a series and integrating, we get

$$\begin{aligned}&\textstyle\sum_{k+j\geq 0} \lambda^{k+j}(t)a^{k+2j}(k!j!)^{-1} \int_0^1 \varphi^k(s_1;t)\rho(x|s_1;t)ds_1 - \\ &\textstyle\sum_{k+j\geq 0} \lambda^{k+j}(t)a^{k+2j}(k!j!)^{-1}\varphi^k(s_1;t)\rho(x|s_1;t) + \\ &\textstyle\sum_{k+j\geq 0} \lambda^{k+j}(t)a^{k+2j+1}(k!j!)^{-1}x(s_1;t) \int_0^1 \varphi^k(s_1;t)\rho(x|s_1;t)ds_1 = 0.\end{aligned}$$

To simplify this equation we extract two first terms corresponding to the indices $k = 0, j = 0$ and $k = 1, j = 0$, so that the remainder of the sum is taken over the set of indices $N(k,j) = \{(k,j) \neq (0,0),(1,0)\}$. After some algebra we obtain the equation

$$\begin{gathered} x(s_1;t) - \lambda(t)\varphi(s_1;t) + \textstyle\sum_{N(k,j)} \lambda^{k+j}(t) a^{k+2j-1}(k!j!)^{-1} \times \\ \times \left[\varphi^k(s_1;t)\rho(x|s_1;t) - (1 + ax(s_1;t)) \textstyle\int_0^1 \varphi^k(s_1;t)\rho(x|s_1;t)ds_1\right] = 0. \end{gathered} \tag{25}$$

Note that for $a = 0$ the “principal” solution of this equation satisfying the normalization condition is $x_0(t,s) = \lambda_0(t)\varphi(s;t)$, where $\lambda_0(t) = (2\sigma^2_\varphi(t))^{-1}$. Our aim is to build the “perturbation” of this solution for $a > 0$.

We introduce a new small functional parameter $\nu(t) = \lambda(t) - \lambda_0(t)$ and a new unknown function $y(t,s) = x(t,s) - \lambda_0(t)\varphi(s;t)$, and substitute them into equation (25). We have

$$\begin{aligned} & y(s_1;t) - \nu(t)\varphi(s_1;t) + \\ & \textstyle\sum_{N(k,j)} (\nu(t) + \lambda_0(t))^{k+j} a^{k+2j-1}(k!j!)^{-1}[\varphi^k(s_1;t)\rho^j(y + \lambda_0(t)\varphi(s_1;t)|s_1;t) - \\ & \textstyle\int_0^1 \varphi^k(s_1;t)\rho^j(y + \lambda_0(t)\varphi(s_1;t)|s_1;t)ds_1] - \sum_{N(k,j)} (\nu(t) + \lambda_0(t))^{k+j} a^{k+2j}(k!j!)^{-1} \times \\ & (y(s_2;t) + \lambda_0(t)\varphi(s_2;t)) \textstyle\int_0^1 \varphi^k(s_1;t)\rho^j(y(s_1,t) + \lambda_0(t)\varphi(s_1;t)|s_1,t)ds_1 = 0. \end{aligned} \tag{26}$$

Due to the inequality

$$\lambda_0(t) = (2\sigma^2_\varphi(t))^{-1} < \varphi_0^{-2}, \quad t \in T,$$

and the boundedness of the kernel, the series on the left-hand side are convergent series of k, j-linear operators (see [20]) in y and ν with bounded coefficients. Therefore the left-hand side of equation (26) is the analytic operator

$$A(y,\nu,a) : B(I \times T) \times B(T) \times R^1 \to B(I \times T).$$

The Fréchet derivative $A_y(y,0,0)$ at the point $y = 0$ is the unit operator and hence is bounded. Then Theorem 3.1 guarantees the existence of a solution of the form

$$y(s_1;t) = \sum_{k+j \geq 1} c_{kj}(s_1;t)\nu^k(t)a^j, \tag{27}$$

where the series is absolutely convergent in the space $B(I \times T)$ for sufficiently small $||\nu||$ and a. This means the convergence of the power series with numerical coefficients

$\sum_{k+j\ge 1}||c_{kj}||\cdot||\nu||^k a^j$ for sufficiently small $||\nu||$ and a. Note that in the proof of Theorem 3.1 in [20] the majorant series for the solution was built explicitly.

Now we substitute solution (27) into equation (26). Equating the coefficients at the same powers of ν and a, we obtain the expressions for c_{kj}. For example,

$$c_{10}(s;t)=\varphi(s;t),\; c_{01}(s;t)=\frac{1}{2}\lambda_0^2(t)(\varphi(s;t)-3\sigma_\varphi^2(t))+\lambda_0(t)\rho(s;t),$$

and so on. Returning to the function x, we have

$$x(s;t)=\lambda_0(t)\varphi(s;t)+\sum_{k+j\ge 1}c_{kj}(s;t)\nu^k(t)a^j. \tag{28}$$

Substituting this solution into normalization condition (24), we obtain

$$\begin{aligned}&2\textstyle\sum_{k+j\ge 1}\int_0^1 c_{kj}(s;t)\varphi(s;t)ds\;\nu^k(t)a^j\\&+\textstyle\sum_{k+j\ge 1}\sum_{i+l\ge 1}\int_0^1\int_0^1\Phi(s_1,s_2;t)c_{kj}(s_1;t)c_{il}(s_2;t)ds_1ds_2\;a^{i+j}\nu^{k+l}(t)=1.\end{aligned} \tag{29}$$

As $c_{10}(s;t)=\varphi(s;t)$, the coefficient at $\nu(t)$ is equal to $2\sigma^2(t)$ and is positive on T. Dividing by $2\sigma^2(t)$, we have the equation

$$\nu(t)=\sum_{k\ge 1,l\ge 2}\gamma_{kl}(t)a^k\nu^l(t), \tag{30}$$

where, as seen from (29), the series with coefficients $\gamma_{kl}(t)$ converges absolutely in some ball of the space $B(T)$.

Applying again Theorem 3.1 to equation (29), we obtain the representation

$$\nu(t)=\sum_{p\ge 1}e_p(t)a^p, \tag{31}$$

where the series converges absolutely in $B(T)$ for sufficiently small $a>0$. Substituting (31) into (28), we get again the convergent series. Returning to the extremal $x(s,t)=\lambda_0(t)x_0(s;t)+y(t)$, we substitute the new series for x into (20). Integrating term-wise and using the convergence of the series for the solution, we obtain the expression for $g(a,t)$ of the form

$$g(a,t|\Phi)=\sum_{i=2}^{\infty}c_i(t)a^i,\; t\in T, \tag{32}$$

where $c_2(t)=(8\sigma_\varphi^2(t))^{-1}$, and the series is convergent for sufficiently small positive a, so that the majorant series $\sum_{i=1}^{\infty}||c_i||a^i$ is also convergent.

Now we can prove the continuity in a and other properties of the function $g_T(a|\Phi)$ listed in Theorem 2.1. Indeed, for any $a_1 > a_2$ from the interval of convergence of the majorant series we have

$$\begin{aligned}|g_T(a_1|\Phi) - g_T(a_2|\Phi)| &\le \sup_{t\in T} |\textstyle\sum_{k=2}^{\infty} c_k(t)(a_1^k - a_2^k)| \\ &\le |a_1 - a_2| \textstyle\sum_{k=2}^{\infty} k \ ||c_k|| a_1^{k-1}.\end{aligned}$$

The series $\sum_{k=2}^{\infty} k \ ||c_k|| a_1^{k-1}$ has the same radius of convergence as the majorant series $\sum_{i=1}^{\infty} ||c_i|| a^i$, hence the sum $\sum_{k=2}^{\infty} k \ ||c_k|| a_1^{k-1}$ is bounded, and the continuity of $g_T(a|\Phi)$ follows.

Now let us estimate the difference $|g_T(a|\Phi) - a^2/(8\varphi_0^2)|$ for small a. Note that

$$g_T(a|\Phi) = \inf_{t\in T} g(t, a|\Phi) \le a^2/8\varphi_0^2 + \sup_{t\in T} |\sum_{k\ge 3} c_k(t) a^k|.$$

On the other hand,

$$g_T(a|\Phi) = \inf_{t\in T} g(t, a|\Phi) \ge a^2/8\varphi_0^2 - \sup_{t\in T} \sum_{k\ge 3} (-c_k(t)) a^k.$$

Hence we obtain

$$|g_T(a|\Phi) - a^2/(8\varphi_0^2)| \le \textstyle\sum_{k=3}^{\infty} ||c_k|| a^k = a^3 \sum_{k=3}^{\infty} ||c_k|| a^{k-3} = O(a^3), \ a \to 0.$$

For kernels of degree $m > 2$ in (12), the term $2m^2\varphi_0^2$ appears instead of $8\varphi_0^2$. □

4 Upper bound.

In this section, we bound the large deviation probabilities from above. For this we apply the exponential inequality for non-degenerate U-statistics from [22, Theorem 2.] For simplicity, we give here a slightly weaker version of it.

Lemma 4.1 *Under the conditions of Theorem 1.2 with $\sigma^2 > 0$ and $z > 0$ we have*

$$\mathbb{P}(|U_n| > z) \le 4 \exp\left(-\frac{nz^2}{2m^2\sigma^2 + Lz}\right),$$

where $L := (2^{m+3} m^m + (2/3) m^{-1})\, M$.

Proof of Theorem 2.2. Consider the partition of the parametric set $a = t_0 < t_1 < \dots < t_N = b$ from the monotonicity condition with the nodes different from the zeros of the variance function $\sigma_\varphi^2(t)$. On any interval of the form $[t_k, t_{k+1}),\ k = 0, \dots, N-1$, we have

$$\sup_{t_k \le t < t_{k+1}} U_n(t) \le U_n(t_{k+1}) + \Delta_n(N). \tag{33}$$

Next, using (33) for $\tau_N > 0$ from condition (15),

$$\begin{aligned} \mathbb{P}(\sup_t U_n(t) > a) &\le \textstyle\sum_{k=0}^{N-1} \mathbb{P}(\sup_{t_k \le t < t_{k+1}} U_n(t) \ge a) \le \\ &\le \textstyle\sum_{k=0}^{N-2} \mathbb{P}(U_n(t_{k+1}) \ge a - \tau_N) + N\mathbb{P}(\Delta_n(N) > \tau_N) = \Gamma_{1,N} + \Gamma_{2,N}. \end{aligned}$$

Let us divide the sum $\Gamma_{1,N}$ into two parts: the first sum includes the indices k for which $t_{k+1} \in T$, while the second sum includes all remaining indices. For $k = 0, \dots, N-2$ the rv $U_n(t_{k+1})$ is a U-statistic with centred and bounded kernel Φ. By Theorem 1.2 we have for the summands of the first sum

$$\mathbb{P}(U_n(t_{k+1}) > a - \tau_N) = \exp(-ng(a - \tau_N, t_{k+1}) + o(n)) \le \exp(-ng_T(a - \tau_N) + o(n)).$$

For the summands of the second sum, by Lemma 4.1 for $t_{k+1} \notin T$

$$\mathbb{P}(U_n(t_{k+1}) > a - \tau_N) \le 4 \exp\left(-\frac{n(a - \tau_N)^2}{m^2\varphi_0^2 + L(a - \tau_N)}\right).$$

Therefore

$$\Gamma_{1,N} \le N \exp(-ng_T(a - \tau_N|\Phi) + o(n)) + 4N \exp\left(-\frac{n(a - \tau_N)^2}{m^2\varphi_0^2 + L(a - \tau_N)}\right). \tag{34}$$

Thanks to (16) the term $\Gamma_{2,N}$ decreases faster than $\Gamma_{1,N}$, and can be neglected. Taking the logarithms in the inequality (34), dividing by n and passing to the limit as $n \to \infty$, we obtain

$$\limsup_{n\to\infty} n^{-1} \ln \mathbb{P}(\sup_t U_n(t) > a) \le -\min\left(g_T(a - \tau_N|\Phi), \frac{(a - \tau_N)^2}{m^2\varphi_0^2 + L(a - \tau_N)}\right).$$

By continuity of the function $g_T(\cdot\ |\Phi)$ as $N \to \infty$

$$\limsup_{n\to\infty} n^{-1} \ln \mathbb{P}(\sup_t U_n(t) > a) \le -\min\left(g_T(a|\Phi), a^2(m^2\varphi_0^2 + La)^{-1}\right).$$

But for small a, by (12), $g_T(a|\Phi) < a^2(m^2\varphi_0^2 + La)^{-1}$. Hence we obtain the required inequality

$$\limsup_{n\to\infty} n^{-1} \ln \mathbb{P}(\sup_t U_n(t) > a) \le -g_T(a|\Phi). \qquad \square$$

Theorem 2.3 follows immediately from Theorems 2.1 and 2.2.

5 Kolmogorov-Smirnov-type statistics

In this section we prove that the Kolmogorov-Smirnov-type statistics $DU_n^{\pm}$ and $SU_n^{\pm}$ satisfy the monotonicity condition, so that Theorem 2.3 is applicable to them.

First, consider the statistic DU_n^+ and assume that the family of kernels $\Theta(\cdot\,;t)$ is non-degenerate and $\theta_0^2 > 0$. Let N be a large number such that $a - N^{-1} > 0$. We divide the parametric set into N parts with nodes $t_k = G^{-1}(k/N), k = 0, ..., N$, where G is from (2). If some interior node coincides with the zero of the variance function, we make a shift of order $O(N^{-2})$. On any interval $[t_k, t_{k+1}), k = 0, ..., N-1$ we have

$$\sup_{t_k \le t < t_{k+1}} (G_n(t) - G(t)) \le G_n(t_{k+1}) - G(t_{k+1}) + O(N^{-1}).$$

Hence the monotonicity condition (15) holds if we take $\Delta_n(N) = O(N^{-1})$, and τ_N to be any sequence tending to zero slower than N^{-1}. The same procedure is applied to the statistic DU_n^-.

The arguments for SU_n^+ are similar. Take the nodes $t_{k,N} = F^{-1}(\frac{k}{N}), k = 0, ..., N$, and shift them, if necessary, as above. Hence for $k = 0, \ldots, N-1$ we have

$$\sup_{t_k \le t < t_{k+1}} [G_n(t) - F_n(t)] \le G_n(t_{k+1}) - F_n(t_{k+1}) + F_n(t_{k+1}) - F_n(t_k).$$

The part of the quantity $\Delta_n(N)$ in the monotonicity condition plays the rv

$$F_n(t_{k+1}) - F_n(t_k) = n^{-1} \sum_{j=1}^{n} I\{t_k \le X_j < t_{k+1}\}.$$

Obviously, the sum on the right-hand side has the binomial distribution with parameters n and $p = 1/N$. The next lemma is proved in [11].

Lemma 5.1 *Let $Bin(n, 1/N)$ be the rv having binomial distribution with parameters n and $1/N$. Then for any $\tau \in (0,1)$ the following inequality holds*

$$\mathbb{P}(Bin(n, 1/N) > n\tau) \le 4^n \exp(-n\tau \ln N).$$

We apply this Lemma with $\tau = \tau_N = (\ln N)^{-1/2}$. For sufficiently large N

$$\mathbb{P}(Bin(n, 1/N) > n/\sqrt{\ln N}) \le \exp(-\frac{1}{2} n \sqrt{\ln N}),$$

and hence

$$\mathbb{P}(F_n(t_{k+1}) - F_n(t_k) > \tau_N) \le \mathbb{P}(Bin(n, 1/N) > n\tau_N) \le \exp(-\frac{1}{2} n \sqrt{\ln N}). \qquad (35)$$

From this the monotonicity condition follows. The arguments for SU_n^- are similar.

We see that Theorem 2.2 is applicable to statistics $DU_n^\pm$ and $SU_n^\pm$ if corresponding families of kernels are non-degenerate and centred. In this case

$$\limsup_{n\to\infty} n^{-1} \ln \mathbb{P}(DU_n^\pm > a) \leq -g_T(a| \pm \Theta),$$

$$\limsup_{n\to\infty} n^{-1} \ln \mathbb{P}(SU_n^\pm > a) \leq -g_T(a| \pm \Psi).$$

Together with (13) and (14) this implies

$$\lim_{n\to\infty} n^{-1} \ln \mathbb{P}(DU_n^\pm > a) = -g_T(a| \pm \Theta),$$

$$\lim_{n\to\infty} n^{-1} \ln \mathbb{P}(SU_n^\pm > a) = -g_T(a| \pm \Psi).$$

From these relations, as explained at the end of section 2, Theorem 2.4 follows.

In order to illustrate the result on large deviations of the statistic DU_n, assume for simplicity that the initial continuous d.f. F is defined on some finite or infinite interval $[a, b]$ and is strictly monotonic there. Consider the kernel $h(x, y) = \max(x, y)$. Then $EI\{h(X, Y) < t\} = F^2(t)$. It follows that the kernels corresponding to one-sided statistics $DU_n^{(max)}$ have the form $\Theta(x, y; t) = I\{\max(x, y) < t\} - F^2(t), t \in R^1$. The projections of these kernels are

$$\theta(x; t) = \mathbb{P}(\max(x, Y) < t) - F^2(t) = I\{x < t\}\ F(t) - F^2(t),$$

and the common variance function is

$$\sigma_\theta^2(t) = E\theta^2(X; t) = F^3(t)(1 - F(t)).$$

The maximum of this function is 27/256. It now follows that for some continuous function v_0

$$\lim_{n\to\infty} n^{-1} \ln \mathbb{P}(DU_n^{(\max)} \geq a) = v_0(a) = -\frac{32}{27}a^2 + O(a^3),\ a \to 0.$$

6 Statistical applications

In this section we apply our general theorems proved above to particular U-empirical tests of Kolmogorov-Smirnov type. At the same time, we fill small gaps in the proofs of [10, 11] and [25], where the incorrect paper [23] was used.

1. *Test of exponentiality based on Desu's characterization.* Let us return to the statistic DE_n given in the Introduction. This statistic is scale-free, so we can

assume that the observations have standard exponential distribution. The kernel of the family of U-statistics DE_n^+ takes the equivalent form

$$\Psi(x,y;t) = \frac{1}{2}(I\{x>t\}+I\{y>t\}) - I\{\min(x,y)>t/2\}, \quad x,y,t\geq 0.$$

The projection is given by

$$\begin{aligned}&\psi(y;t) = \tfrac{1}{2}(\mathbb{P}\{X_1>t\}+I\{y>t\}] - E[I\{\min(X_1,y)>t/2\}]\\ &= \tfrac{1}{2}\exp(-t)+\tfrac{1}{2}I\{y>t\} - \exp(-t/2)\ I\{y>t/2\},\end{aligned}$$

and hence the variance function $\sigma_\psi^2(t) = \frac{1}{4}\exp(-t)(1-\exp(-t)), t\geq 0$. So, the family of kernels is non-degenerate, and we can apply Theorem 2.4. It is seen that $\psi_0^2 = \sup_{t\geq 0}\sigma_\psi^2(t) = \frac{1}{16}$, and this determines the first term of the large deviation asymptotics. So, there exists a continuous function v_1 such that

$$\lim_{n\to\infty} n^{-1}\ln\mathbb{P}(DE_n>a) = v_1(a) = -2a^2+O(a^3), \quad a\to 0. \tag{36}$$

2. Another test of exponentiality is based on the simplified "lack of memory" property, see [8] and [11]. Consider the statistic

$$AN_n^+ = \sup_{x\geq 0}[\bar{F}_n(2x) - \bar{F}_n^2(x)].$$

Statistics AN_n^- and AN_n are defined analogously. Large values of these statistics are statistically significant.

It was shown in [11] that under the hypothesis of exponentiality the statistic AN_n^+ admits the representation

$$AN_n^+ = \sup_{0<t<1}\binom{n}{2}^{-1}\sum_{1\leq i<j\leq n}\Psi(U_i,U_j;t)+O(n^{-1}),$$

where $U_1,...,U_n$ are uniformly distributed on [0,1] rv's. The family of kernels is then given by

$$\begin{aligned}\Psi(x_1,x_2;t) = I\{x_1<t\}+I\{x_2<t\}-I\{x_1<t\}I\{x_2<t\}-\\ (1/2)(I\{x_1<2t-t^2\}+I\{x_2<2t-t^2\}).\end{aligned}$$

The corresponding family of U-statistics satisfies the monotonicity condition [11]. Simple calculations show that

$$\psi(z;t) = E\left(\Psi(X_1,X_2;t)|X_1=z\right) = (1-t)I\{z<t\}+t^2/2-(1/2)I\{z<2t-t^2\}$$

and hence the variance function is equal to

$$\sigma_\psi^2(t) = (1/4)t(1-t)^2(2-t) > 0, \quad 0 < t < 1.$$

We can apply Theorem 2.4.

The maximum of the variance function is attained at $t = 1 - 1/\sqrt{2}$ and is equal to $1/16$. Therefore for some continuous function v_2

$$\lim_{n\to\infty} n^{-1} \ln \mathbb{P}(AN_n^+ \geq a) = v_2(a) = -2a^2 + O(a^3),\ a \to 0.$$

The same asymptotics is valid for statistics AN_n^- and AN_n.

3. One more characterization of the exponential law belongs to Puri and Rubin [24]: *Let X and Y be independent rv's with common absolutely continuous df F on R^+. F is exponential if and only if $|X-Y|$ has the same distribution as X.*

We construct a one-sided U-empirical version of the Kolmogorov-Smirnov test by introducing the statistic

$$PR_n^+ = \sup_{t\geq 0} \left(\binom{n}{2}^{-1} \sum_{1\leq i<j\leq n} I\{|X_i - X_j| < t\} - n^{-1}\sum_{k=1}^n I\{X_k < t\} \right).$$

Statistics PR_n^- and PR_n are defined analogously. Thus we obtain the family of U-statistics with the kernels

$$\Psi(x_1, x_2; t) = I\{|x_1 - x_2| < t\} - \frac{1}{2}(I\{x_1 < t\} + I\{x_2 < t\}),\ t \geq 0.$$

Hence

$$\psi(s;t) = E\left(I\{|X_1 - s| < t\} - \tfrac{1}{2}(I\{X_1 < t\} + I\{s < t\})\right) = \\ = \left(e^{-s+t} - \frac{1}{2}\right) I\{s \geq t\} - e^{-t}(e^{-s} - \frac{1}{2}),$$

and the variance function is equal to

$$\sigma_\psi^2(t) = \frac{1}{12} e^{-t}(1 + e^{-t} - 2e^{-2t}),\ t \geq 0.$$

The maximum of this function is attained for $e^{-t} = \frac{\sqrt{7}+1}{6}$ and is equal to $\frac{10+7\sqrt{7}}{648}$. Hence there exists a continuous function v_3 such that as $a \to 0$

$$\lim_{n\to\infty} n^{-1} \ln \mathbb{P}(PR_n > a) = v_3(a) = -\frac{7\sqrt{7}-10}{3} a^2 + O(a^3) \approx -2.840a^2 + O(a^3).$$

The next two examples are related to the problem of testing symmetry.

4. Let $X_1, X_2, \ldots, X_n$ be a random sample from a continuous df F_0 . We wish to test the hypothesis of symmetry about zero

$$H_0 : \Delta F_0(x) := F_0(x) + F_0(-x) - 1 = 0 \quad \forall x \in \mathbb{R}^1.$$

Consider the centered (in time) Kolmogorov-Smirnov statistics [25]. One of the one-sided statistics has the form

$$H_n^+ = \sup_t \left[\Delta F_n(t) - \int_{-\infty}^{\infty} (\Delta F_n(y))\, dF_n(y) \right],$$

the second statistic H_n^- is introduced in a similar manner, and $H_n = \max(H_n^+, H_n^-)$.

Formally, these statistics do not belong to the class of statistics $SU_n^{\pm}$ and SU_n, however the difference between them is nonsignificant. Without loss of generality we can assume that the distribution of X_i is uniform on $[-1, 1]$. For any $t \in [-1, 1]$ consider the family of statistics

$$H_n^+(t) = F_n(t) + F_n(-t) - \int_{-1}^{1} (F_n(t) + F_n(-t)) dF_n(t) = n^{-2} \sum_{i,j=1}^{n} \Psi(X_i, X_j; t) + n^{-1},$$

where the family of kernels $\Psi(\cdot\,; t)$ has the form

$$\begin{aligned} \Psi(x_1, x_2; t) = \frac{1}{2} \Big(& I\{x_1 < t\} + I\{x_1 < -t\} + I\{x_2 < t\} + I\{x_2 < -t\} \\ & - 2 \cdot I\{x_1 + x_2 < 0\} - 1 \Big). \end{aligned}$$

Now we introduce the auxiliary family of statistics

$$\mathcal{H}_n^+(t) = \binom{n}{2}^{-1} \sum_{1 \le i < j \le n} \Psi(X_i, X_j; t).$$

Note that for any t and any $n > 1$

$$|H_n^+(t) - \mathcal{H}_n^+(t)| \le 7n^{-1},$$

so the large deviation asymptotics for $\sup_t H_n^+(t)$ and $\sup_t \mathcal{H}_n^+(t)$ are the same. It is easy to check that

$$\psi(z; t) \equiv E\left(\Psi(X_1, X_2; t) | X_1 = z\right) = \frac{1}{2} \left(I\{z < t\} + I\{z < -t\} + t - 1 \right),$$

and hence

$$\sigma_\psi^2(t) = \frac{1}{4}(t^2 - |t| + \frac{1}{3}), \ -1 \le t \le 1.$$

Clearly $\psi_0^2 = \sup_{-1 \le t \le 1} \sigma_\psi^2(t) = 1/12$.

Let us turn to the upper bound and consider the uniform partition of $[-1, 1]$ into $2N$ parts using the nodes $t_k = k/N,\ k = -N, ..., N$. Obviously,

$$\sup_{t_k \le t < t_{k+1}} \mathcal{H}_n^+(t) \le \mathcal{H}_n^+(t_{k+1}) + F_n(t_{k+1}) - F_n(t_k).$$

By Lemma 5.1 this ensures the monotonicity condition. Therefore for some continuous function v_4 as $a \to 0$

$$\lim_{n\to\infty} n^{-1} \ln \mathbb{P}(H_n^+ > a) = v_4(a) = -\frac{3}{2}a^2 + O(a^3).$$

Similar results hold for H_n^- and H_n.

5. Another test of symmetry is based on the characterization established by Baringhaus and Henze [9]: *The common distribution of two independent rv's X and Y is symmetric with respect to zero iff $|X|$ and $|\max(X, Y)|$ have the same distribution.*

Consider two edf's based on the sample $X_1, ..., X_n$. Let

$$L_n(x) = n^{-1} \sum_{j=1}^{n} I(|X_j| \le x), x \ge 0$$

and

$$G_n(x) = \binom{n}{2}^{-1} \sum_{1 \le j < k \le n} I(|\max(X_j, X_k)| \le x), x \ge 0.$$

Following [9], consider the statistic

$$BH_n^+ = \sup_{x \ge 0}[L_n(x) - G_n(x)].$$

The statistics BH_n^- and BH_n are defined analogously. All these statistics are distribution-free, and we may assume that the observations are uniformly distributed on $[-1, 1]$. The statistic BH_n^+ admits the representation

$$BH_n^+ = \binom{n}{2}^{-1} \sup_{0<t<1} \sum_{1 \le j < k \le n} \Xi(X_j, X_k; t),$$

where for any $t \in (0, 1)$

$$\Xi(X_j, X_k; t) = I\{|\max(X_j, X_k)| \le t\} - \frac{1}{2}(I\{|X_j| \le t\} + I\{|X_k| \le t\}).$$

Formally this family does not fit our theory, however replacing edf F_n by edf L_n leads to minimal changes in the proofs. Simple calculations show that for any $z \in [-1,1]$ and $0 < t < 1$ the projections of the kernels have the form

$$\xi(z;t) = E\left(\Xi(X_1, X_2; t) | X_1 = z\right) = \begin{cases} t/2, & -1 \le z < -t \\ 0, & -t \le z \le t \\ -t/2, & t < z \le 1. \end{cases}$$

Consequently the variance function is given by

$$\sigma_\xi^2(t) = \frac{t^2}{8} \int_{-1}^{1} (I\{-1 \le x < -t\} - I\{t < x \le 1\})^2 dx = \frac{1}{4} t^2 (1 - |t|), -1 \le t \le 1.$$

The maximum of this function is attained at $\pm\frac{2}{3}$ and is equal to $\frac{1}{27}$. Hence the large deviation asymptotics has the form

$$\lim_{n\to\infty} n^{-1} \ln \mathbb{P}(BH_n^+ \ge a) = v_5(a) = -\frac{27}{8} a^2 + O(a^3),\ a \to 0.$$

Similar statements hold true for the statistics BH_n^- and BH_n.

6. Consider the famous characterization of normality due to G. Polya [26]: *Let X and Y be i.i.d. rv's with zero mean. Then X and $(X+Y)/\sqrt{2}$ have the same distribution iff X and Y are normally distributed with some positive variance.*

The integral test of normality based on this characterization was proposed in [27]. Let us construct the scale-invariant Kolmogorov-type test comparing the usual edf and the U-empirical df, based on $\frac{X_i + X_j}{\sqrt{2}}$. We arrive at the one-sided statistic

$$PO_n^+ = \sup_t \left(\binom{n}{2}^{-1} \sum_{1 \le i < j \le n} I\{X_i + X_j < t\sqrt{2}\} - n^{-1} \sum_{k=1}^{n} I\{X_k < t\} \right).$$

The statistics PO_n^- and PO_n are introduced similarly. Our statistic PO_n^+ corresponds to the family of kernels

$$\Psi(x_1, x_2; t) = I\{x_1 + x_2 < t\sqrt{2}\} - \frac{1}{2}(I\{x_1 < t\} + I\{x_2 < t\})$$

and satisfies the monotonicity condition.

Denote by $\mathcal{N}$ the df of the standard normal law. Then the projections of the kernels are

$$\psi(s;t) = E(\Psi(X_1, X_2; t) | X_2 = s) = \mathcal{N}(t\sqrt{2} - s) - \frac{1}{2}\mathcal{N}(t) - \frac{1}{2} 1\{s < t\}.$$

Consequently, the variance function is

$$\sigma_{\psi}^{2}(t)=\int_{-\infty}^{\infty}\mathcal{N}^{2}(t\sqrt{2}-s)d\mathcal{N}(s)-\int_{-\infty}^{t}\mathcal{N}(t\sqrt{2}-s)d\mathcal{N}(s)+\frac{1}{4}\mathcal{N}(t)-\frac{1}{4}\mathcal{N}^{2}(t).$$

The problem of finding the maximum of this function analytically is difficult. However, its plot clearly shows that this maximum is attained at zero, and consequently is equal to 1/48.



Fig.1 *Plot of the variance function for Polya test.*

Therefore there exists a continuous function v_6 such that

$$\lim_{n\to\infty} n^{-1}\ln\mathbb{P}(PO_n>a)=v_6(a)=-6a^2+O(a^3),\ a\to 0.$$

7 Calculation of local Bahadur efficiency

The results on large deviations allow us to calculate local Bahadur efficiency of U-empirical Kolmogorov-Smirnov tests.

As an illustration we find the efficiency of Desu's test for some parametric alternatives F_θ to the hypothesis of exponentiality. The local exact slope, see [3], is the main part as $\theta\to 0$ of the expression

$$4\sup_{x}|(1-F_\theta(x/2))^2-(1-F_\theta(x))|^2.$$

Take, for example, the Weibull alternative with $1 - F_\theta(x) = \exp(-x^{1+\theta}), \theta \geq 0$. Then the local exact slope is equivalent to

$$4 \ln^2(2) \sup_x (x \exp(-x))^2 \ \theta^2 \approx 0.2601 \ \theta^2.$$

The theoretical maximum (double Kullback-Leibler information, see [2]), is equal [11] to $\pi^2/6 \cdot \theta^2 \approx 1.6449 \ \theta^2$. Hence the local Bahadur efficiency of Desu's test equals 0.1581. At the same time the efficiency of Desu's test for Makeham alternative with the density $(1 + \theta(1 - e^{-x})) \exp(-x - \theta[x - (1 - e^{-x})]), x \geq 0, \theta \geq 0$, is much larger and is equal to 0.4938.

Other calculations of local Bahadur efficiency can be found in [25], [10] and [11]. It turns out that in some cases U-empirical Kolmogorov tests have high efficiency and perform well compared to some other goodness-of-fit tests. For instance, the local efficiency of the sequence of statistics H_n for testing symmetry under the normal shift alternative is equal to 0.955. In the same problem the local efficiency of the sequence of statistics BH_n is equal to 0.75.

It would be interesting to construct new U-empirical tests of Kolmogorov-Smirnov type and to calculate their efficiencies using the large deviation results obtained in this paper. We hope to return to this question later.

Acknowledgements

The author is thankful to Prof. A.I. Nazarov for reading the paper and making useful comments and to Prof. R. Serfling for valuable advice.

References

[1] Abrahamson I. G. *Exact Bahadur efficiencies for the Kolmogorov - Smirnov and Kuiper one- and two-sample statistics.* Ann. Math. Stat., 38(1967), pp. 1475–1490.

[2] Bahadur R. R. *Some limit theorems in statistics*, SIAM, Philadelphia, 1971.

[3] Nikitin Y. *Asymptotic efficiency of nonparametric tests.* Cambridge University Press, 1995.

[4] Serfling R. J. *Generalized L-, M-, and R-statistics.* Ann. Stat., 12(1984), pp. 76-86.

[5] Helmers R., Janssen P., Serfling R. *Glivenko-Cantelli properties of some generalized empirical DF's and strong convergence of generalized L-statistics.* Probab. Theory Relat. Fields 79(1988), pp. 75-93.

[6] Janssen P.L. *Generalized empirical distribution functions with statistical applications*, Limburgs Universitair Centrum, Diepenbeek, 1988.

[7] Serfling R.J., Wang W. *A large deviation theorem for U-processes.* Stat. Probab. Lett. 49(2000), pp. 181-193.

[8] Angus J. E. *Goodness-of-fit tests for exponentiality based on a loss-of-memory type functional equation,* J. Statist. Planning Infer., 6(1982), pp. 241-251.

[9] Baringhaus L., Henze N., *A characterization of and new consistent tests of symmetry*, Commun. Statist.- Theor. Meth., 21(1992), pp. 1555-1566.

[10] Nikitin Ya. Yu. *On Baringhaus-Henze test for symmetry: Bahadur efficiency and local optimality for shift alternatives.* Math. Methods of Statist., 5(1996), pp. 214-226.

[11] Nikitin Ya. Yu. *Bahadur efficiency of a test of exponentiality based on a loss-of-memory type functional equation.* J. of Nonparam. Statist., 6(1996), pp. 13 - 26.

[12] Desu M.M. *A characterization of the exponential distribution by order statistics.* Ann. Math. Stat., 42(1971), pp. 837 - 838.

[13] Silverman B. W. *Convergence of a class of empirical distribution functions of dependent random variables.* Ann. Probab. 11(1983), pp. 745-751.

[14] Dehling H., Denker M., Philipp W. *The almost sure invariance principle for the empirical process of U-statistic structure.* Annales de l'I.H.P., B23(1987), 121 - 134.

[15] Serfling R. *Approximation Theorems of Mathematical Statistics.* Wiley, New York, 1980.

[16] Eichelsbacher P., Löwe M. *A Large Deviation Principle for m-variate von Mises-statistics and U-Statistics.* J. Theoret. Prob., 8(1995), pp. 807–823.

[17] Arcones M. A. *Large deviations for U-statistics.* Journ. of Multiv. Anal., 42(1992), pp. 299-301.

[18] Nikitin Ya. Yu., Ponikarov E. V. *Rough large deviation asymptotics of Chernoff type for von Mises functionals and U-statistics.* Proceedings of St.Petersburg Mathematical Society, 7(1999), pp. 124–167. English translation in AMS Transl., ser.2, 203(2001), pp. 107 - 146.

[19] Baringhaus, L., Rank, R. *On large deviations of U-statistics and their projections.* Sankhyā, A64(2002), pp. 167 - 170.

[20] Vainberg M. M., Trenogin V. A. *Theory of Branching of Solutions of Nonlinear Equations,* Noordhoof, Leiden, 1974.

[21] Krasnoselskii M.A., Vainikko V.M., Zabreiko P.V., Rutitskii Ya.B., Stezenko V.A. *Approximate solutions of operator equations.* Wolters-Noordhoff, Groningen, 1972.

[22] Arcones M. A. *A Bernstein-type inequality for U-statistics and U-processes.* Statistics and Prob. Letters, 22(1995), pp. 239 -247.

[23] Dasgupta R. *On large deviation probabilities of U-statistics in non-i.i.d. case.* Sankhyā, 46(1984), pp. 110–116.

[24] Puri P. S., Rubin H. *A characterization based on the absolute difference of two i.i.d. random variables.* Ann. Math. Stat. 41(1970), pp. 2113-2122.

[25] Abbakumov V. L., Nikitin Ya. Yu. *Bahadur efficiency and local optimality of a new nonparametric test of symmetry*, In: Statistics and Probability. A R.R. Bahadur Festschrift, Wiley Eastern Ltd., New Delhi, 1993, pp. 1-12.

[26] Polya G. *Herleitung des Gauss'schen Fehlergesetzes aus einer Funktionalsgleichung.* Math. Zeitschrift, 18(1923), pp. 96 – 108.

[27] Muliere P., Nikitin Ya. *Scale-invariant test of normality based on Polya's characterization.* Metron, 60(2002), N 1-2, pp. 21 – 33.

Department of Mathematics and Mechanics,
St.Petersburg State University,
Universitetski pr. 28, Stary Peterhof,
198504, Russia

e-mail yanikit47@gmail.com